

Research Article

Monire Hajmohamadi, Rahmatollah Lashkaripour* and Mojtaba Bakherad

Further refinements of generalized numerical radius inequalities for Hilbert space operators

<https://doi.org/10.1515/gmj-2019-2023>

Received September 29, 2016; accepted May 21, 2018

Abstract: In this paper, we show some refinements of generalized numerical radius inequalities involving the Young and Heinz inequality. In particular, we present

$$w_p^p(A_1^* T_1 B_1, \dots, A_n^* T_n B_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n [B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp} \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x),$$

where $T_i, A_i, B_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$), f and g are nonnegative continuous functions on $[0, \infty)$ satisfying $f(t)g(t) = t$ for all $t \in [0, \infty)$, $p, r \geq 1$, $N \in \mathbb{N}$, and

$$\begin{aligned} \eta(x) = & \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j}} \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j} \right. \\ & \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Keywords: Euclidean operator radius, Heinz means, numerical radius, positive operator, Young inequality

MSC 2010: Primary 47A12; secondary 47A63, 47A30

1 Introduction

Let $\mathbb{B}(\mathcal{H})$ denote the C^* -algebra of all bounded linear operators on a complex Hilbert space \mathcal{H} with an inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $\|\cdot\|$. In the case where $\dim \mathcal{H} = n$, we identify $\mathbb{B}(\mathcal{H})$ with the matrix algebra \mathbb{M}_n of all $n \times n$ matrices with entries in the complex field. The numerical radius of $T \in \mathbb{B}(\mathcal{H})$ is defined by

$$w(T) := \sup\{|\langle Tx, x \rangle| : x \in \mathcal{H}, \|x\| = 1\}.$$

It is well known that $w(\cdot)$ defines a norm on $\mathbb{B}(\mathcal{H})$, which is equivalent to the usual operator norm $\|\cdot\|$. In fact, for any $T \in \mathbb{B}(\mathcal{H})$, $\frac{1}{2}\|T\| \leq w(T) \leq \|T\|$, see [4]. The quantity $w(T)$ is useful in studying perturbation, convergence and approximation problems as well as interactive methods, etc. For more information, see [1, 3, 5, 6, 11, 17, 18].

The classical Young inequality says that if $0 \leq v \leq 1$, then $a^v b^{1-v} \leq va + (1-v)b$ ($a, b > 0$). During the last decades several generalizations, reverses, refinements and applications of the Young inequality in various settings have been given; see [2, 10] and the references therein. A refinement of the scalar Young inequality is presented in [10] as follows:

$$a^v b^{1-v} \leq va + (1-v)b - r_0(a^{\frac{1}{2}} - b^{\frac{1}{2}})^2, \quad r_0 = \min\{v, 1-v\}. \quad (1.1)$$

*Corresponding author: Rahmatollah Lashkaripour, Department of Mathematics, Faculty of Mathematics, University of Sistan and Baluchestan, Zahedan, Iran, e-mail: mojtaba.bakherad@yahoo.com, bakherad@member.ams.org

Monire Hajmohamadi, Mojtaba Bakherad, Department of Mathematics, Faculty of Mathematics, University of Sistan and Baluchestan, Zahedan, Iran, e-mail: monire.hajmohamadi@yahoo.com, lashkari@hamoon.usb.ac.ir

Recently, Sababheh and Choi obtained, in [14], a refinement of the Young inequality, that is,

$$a^\nu b^{1-\nu} \leq \nu a + (1-\nu)b - S_N(\nu), \quad (1.2)$$

in which

$$S_N(\nu) := \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \nu + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \left(\sqrt[2^j]{b^{2^{j-1}-k_j} a^{k_j}} - \sqrt[2^j]{a^{k_j+1} b^{2^{j-1}-k_j-1}} \right)^2,$$

where $N \in \mathbb{N}$, $r_j = [2^j \nu]$ and $k_j = [2^{j-1} \nu]$. Here $[x]$ is the greatest integer less than or equal to x . When $N = 1$, inequality (1.2) reduces to (1.1).

It follows from $\nu a + (1-\nu)b \leq (\nu a^r + (1-\nu)b^r)^{\frac{1}{r}}$ ($r \geq 1$) and inequality (1.1) that

$$a^\nu b^{1-\nu} \leq (\nu a^r + (1-\nu)b^r)^{\frac{1}{r}} - S_N(\nu).$$

In particular, for $\nu = \frac{1}{2}$, we get

$$a^{\frac{1}{2}} b^{\frac{1}{2}} \leq \left(\frac{1}{2} \right)^{\frac{1}{r}} (a^r + b^r)^{\frac{1}{r}} - \frac{1}{2} \sum_{j=1}^N \left(\sqrt[2^j]{b^{2^{j-1}-k_j} a^{k_j}} - \sqrt[2^j]{a^{k_j+1} b^{2^{j-1}-k_j-1}} \right)^2.$$

If $N = 1$, then we reach [10, inequality (2.1)] as follows:

$$a^{\frac{1}{2}} b^{\frac{1}{2}} \leq \left(\frac{1}{2} \right)^{\frac{1}{r}} (a^r + b^r)^{\frac{1}{r}} - \frac{1}{2} (a^{\frac{1}{2}} - b^{\frac{1}{2}})^2.$$

Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). The Euclidean operator radius of T_1, \dots, T_n is defined in [13] by

$$w_e(T_1, \dots, T_n) := \sup_{\|x\|=1} \left(\sum_{i=1}^n |\langle T_i x, x \rangle|^2 \right)^{\frac{1}{2}}.$$

In [12], the functional w_p of operators T_1, \dots, T_n for $p \geq 1$ is defined by

$$w_p(T_1, \dots, T_n) := \sup_{\|x\|=1} \left(\sum_{i=1}^n |\langle T_i x, x \rangle|^p \right)^{\frac{1}{p}}.$$

Let $T_1, \dots, T_n \in \mathbb{B}(\mathcal{H})$. Recently, Sheikhhosseini et al. showed in [16] that

$$w_p^p(A_1^* T_1 B_1, \dots, A_n^* T_n B_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n [B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp} \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \zeta(x), \quad (1.3)$$

where

$$\zeta(x) = \frac{1}{2} \sum_{i=1}^n (\langle [B_i^* f^2(|T_i|) B_i]^p x, x \rangle^{\frac{1}{2}} - \langle [A_i^* g^2(|T_i^*|) A_i]^p x, x \rangle^{\frac{1}{2}})^2.$$

They also presented the following inequality:

$$w_p(T_1, \dots, T_n) \leq \frac{1}{2} \left[\sum_{i=1}^n (\| |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} \| - 2 \inf_{\|x\|=1} \zeta_i(x))^p \right]^{\frac{1}{p}}, \quad (1.4)$$

in which, $0 \leq \alpha \leq 1$, $p \geq 1$ and $\zeta_i(x) = \frac{1}{2} (\langle |T_i|^{2\alpha} x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)} x, x \rangle^{\frac{1}{2}})^2$.

In the same paper, they showed

$$w_p^p(T_1, \dots, T_n) \leq \frac{1}{2} \left\| \sum_{i=1}^n (|T_i|^{2\alpha p} + |T_i^*|^{2(1-\alpha)p}) \right\| - \inf_{\|x\|=1} \zeta(x), \quad (1.5)$$

where $\zeta(x) = \frac{1}{2} \sum_{i=1}^n (\langle |T_i|^{2\alpha p} x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{\frac{1}{2}})^2$.

Moreover, they established the inequalities

$$w_p^p(T_1, \dots, T_n) \leq \left\| \sum_{i=1}^n \alpha |T_i|^p + (1-\alpha) |T_i^*|^p \right\| - \inf_{\|x\|=1} \zeta(x), \quad (1.6)$$

and

$$w_p^r(|T_1|, \dots, |T_n|) w_q^r(|T_1^*|, \dots, |T_n^*|) \leq \frac{r}{p} \left\| \sum_{i=1}^n |T_i|^p \right\| + \frac{r}{q} \left\| \sum_{i=1}^n |T_i^*|^q \right\| - \inf_{\|x\|=\|y\|=1} \delta(x, y), \quad (1.7)$$

where $\zeta(x) = \min\{\alpha, 1 - \alpha\} \sum_{i=1}^n (\langle |T_i|^p x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^p x, x \rangle^{\frac{1}{2}})^2$ and

$$\delta(x, y) = \frac{r}{p} \left(\sqrt[2]{\sum_{i=1}^n \langle |T_i| x, y \rangle^p} - \sqrt[2]{\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q} \right)^2.$$

Assume that $X \in \mathbb{B}(\mathcal{H})$. The mixed Heinz means are defined by

$$H_\alpha(A, B) = \frac{A^\alpha X B^{1-\alpha} + A^{1-\alpha} X B^\alpha}{2},$$

in which, $0 \leq \alpha \leq 1$ and $A, B \geq 0$, see [8]. In [15], Sattari et al. showed that

$$w^r(A^\alpha X B^{1-\alpha}) \leq \|X\|^r \|\alpha A^r + (1 - \alpha) B^r\|, \quad (1.8)$$

where $A, B, X \in \mathbb{B}(\mathcal{H})$, with A, B positive, $r \geq 2$ and $0 \leq \alpha \leq 1$.

Using inequality (1.8), they presented an upper bound for the Heinz means of matrices as follows:

$$w^r(H_\alpha(A, B)) \leq \|X\|^r \left\| \frac{A^r + B^r}{2} \right\|. \quad (1.9)$$

In the present paper, we refine inequalities (1.3)–(1.9). We also find an upper bound for the functional w_p .

2 Main results

To prove our numerical radius inequalities, we need several known lemmas. The first lemma is a simple result of the classical Jensen, Young and a generalized mixed Cauchy–Schwarz inequalities [7, 9].

Lemma 2.1. *Let $a, b \geq 0$, $0 \leq v \leq 1$ and $r \neq 0$. Then the following hold:*

- (a) $a^v b^{1-v} \leq v a + (1 - v) b \leq (v a^r + (1 - v) b^r)^{\frac{1}{r}}$ for $r \geq 1$.
- (b) If $T \in \mathbb{B}(\mathcal{H})$ and $x, y \in \mathcal{H}$ are any vectors, then

$$|\langle T x, y \rangle|^2 \leq \langle |T|^{2v} x, x \rangle \langle |T^*|^{2(1-v)} y, y \rangle.$$

- (c) If f, g are nonnegative continuous functions on $[0, \infty)$ which satisfy the relation $f(t)g(t) = t$ ($t \in [0, \infty)$), then

$$|\langle T x, y \rangle| \leq \|f(|T|)x\| \|g(|T^*|)x\|$$

for all $x, y \in \mathcal{H}$.

Lemma 2.2 (McCarty inequality [9]). *Let $T \in \mathbb{B}(\mathcal{H})$, $T \geq 0$, and let $x \in \mathcal{H}$ be a unit vector. Then*

- (a) $\langle T x, x \rangle^r \leq \langle T^r x, x \rangle$ for $r \geq 1$,
- (b) $\langle T^r x, x \rangle \leq \langle T x, x \rangle^r$ for $0 < r \leq 1$.

Now, by using inequality (1.2), we get the first result.

Theorem 2.3. *Let $A, B, X \in \mathbb{B}(\mathcal{H})$, with A, B positive, $r \geq 2$ and $0 \leq v \leq 1$. Then*

$$w^r(A^v X B^{1-v}) \leq \|X\|^r \left[\|v A^r + (1 - v) B^r\| - \inf_{\|x\|=1} \eta(x) \right], \quad (2.1)$$

where

$$\begin{aligned} \eta(x) &= \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} v + (-1)^{r_j+1} \left[\frac{r_j + 1}{2} \right] \right) \\ &\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Proof. Let $x \in \mathcal{H}$ be a unit vector. Then, by Lemma 2.2 and inequality (1.2),

$$\begin{aligned}
|\langle A^v X B^{1-v} x, x \rangle|^r &= |\langle X B^{1-v} x, A^v x \rangle|^r \\
&\leq \|X\|^r \|B^{1-v} x\|^r \|A^v x\|^r \\
&= \|X\|^r \langle B^{2(1-v)} x, x \rangle^{\frac{r}{2}} \langle A^{2v} x, x \rangle^{\frac{r}{2}} \\
&\leq \|X\|^r \langle A^r x, x \rangle^v \langle B^r x, x \rangle^{1-v} \\
&\leq \|X\|^r [v \langle A^r x, x \rangle + (1-v) \langle B^r x, x \rangle] - \|X\|^r \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} v + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
&\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.
\end{aligned}$$

Taking the supremum over $x \in \mathcal{H}$, with $\|x\| = 1$, in the above inequality we deduce the desired inequality. \square

Remark 2.4. Let $N = 1$ in inequality (2.1). Then

$$w^r(A^v X B^{1-v}) \leq \|X\|^r \left[\|v A^r + (1-v) B^r\| - \inf_{\|x\|=1} \eta(x) \right], \quad (2.2)$$

in which, $\eta(x) = r_0 (\langle A^r x, x \rangle^{\frac{1}{2}} - \langle B^r x, x \rangle^{\frac{1}{2}})^2$ and $r_0 = \min\{v, 1-v\}$. Hence, inequality (2.2) is a refinement of inequality (1.8).

Using Theorem 2.3 we can find an upper bound for Heinz means of matrices that is a refinement of (1.9).

Theorem 2.5. Suppose $A, B, X \in \mathbb{B}(\mathcal{H})$, with A, B positive. Then

$$w^r(H_v(A, B)) \leq \|X\|^r \left[\left\| \frac{A^r + B^r}{2} \right\| - \frac{1}{2} \inf \zeta(x) \right],$$

where $r \geq 2$, $0 \leq v \leq 1$, $n \in \mathbb{N}$, and

$$\begin{aligned}
\zeta(x) &= \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
&\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.
\end{aligned}$$

Proof. For a unit vector $x \in \mathcal{H}$, we have

$$\begin{aligned}
\left| \left\langle \frac{A^v X B^{1-v} + A^{1-v} X B^v}{2} x, x \right\rangle \right|^r &\leq \left(\frac{|\langle A^v X B^{1-v} x, x \rangle| + |\langle A^{1-v} X B^v x, x \rangle|}{2} \right)^r \\
&\leq \frac{|\langle A^v X B^{1-v} x, x \rangle|^r + |\langle A^{1-v} X B^v x, x \rangle|^r}{2} \\
&\leq \frac{\|X\|^r}{2} [\langle v A^r + (1-v) B^r x, x \rangle] - \frac{\|X\|^r}{2} \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} v + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
&\quad \times \left(\sqrt[2^j]{b^{2^{j-1}-k_j} a^{k_j}} - \sqrt[2^j]{a^{k_j+1} b^{2^{j-1}-k_j-1}} \right)^2 \\
&\quad + \frac{\|X\|^r}{2} [\langle (1-v) A^r + v B^r x, x \rangle] - \frac{\|X\|^r}{2} \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} v + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
&\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \\
&= \|X\|^r \left[\left\langle \frac{A^r + B^r}{2} x, x \right\rangle \right] - \frac{\|X\|^r}{2} \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
&\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.
\end{aligned}$$

If we take the supremum over $x \in \mathcal{H}$, with $\|x\| = 1$, then we deduce the desired inequality. \square

In the next theorem we show a refinement of inequality (1.3).

Theorem 2.6. *Let $T_i, A_i, B_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$) and let f and g be nonnegative continuous functions on $[0, \infty)$ satisfying $f(t)g(t) = t$ for all $t \in [0, \infty)$. Then*

$$w_p^p(A_1^* T_1 B_1, \dots, A_n^* T_n B_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n [B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp} \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x),$$

where $p, r \geq 1, N \in \mathbb{N}$, and

$$\begin{aligned} \eta(x) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j} \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j}} \right. \\ &\quad \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Proof. Let $x \in \mathcal{H}$ be any unit vector. Then, by Lemma (2.1) (c), Lemma (2.2) (a) and (1.2),

$$\begin{aligned} \sum_{i=1}^n |\langle A_i^* T_i B_i x, x \rangle|^p &= \sum_{i=1}^n |\langle T_i B_i x, A_i x \rangle|^p \\ &\leq \sum_{i=1}^n \|f(|T_i|) B_i x\|^p \|g(|T_i^*|) A_i x\|^p \\ &= \sum_{i=1}^n \langle f(|T_i|) B_i x, f(|T_i|) B_i x \rangle^{\frac{p}{2}} \langle g(|T_i^*|) A_i x, g(|T_i^*|) A_i x \rangle^{\frac{p}{2}} \\ &= \sum_{i=1}^n \langle B_i^* f^2(|T_i|) B_i x, x \rangle^{\frac{p}{2}} \langle A_i^* g^2(|T_i^*|) A_i x, x \rangle^{\frac{p}{2}} \\ &\leq \sum_{i=1}^n \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{\frac{1}{2}} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{\frac{1}{2}} \\ &\leq \sum_{i=1}^n \left[\left(\frac{1}{2} \langle (B_i^* f^2(|T_i|) B_i)^{pr} x, x \rangle + \frac{1}{2} \langle (A_i^* g^2(|T_i^*|) A_i)^{pr} x, x \rangle \right)^{\frac{1}{r}} \right] \\ &\quad - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j} \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j}} \right. \\ &\quad \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \\ &\leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\langle \left(\sum_{i=1}^n ([B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp}) \right) x, x \right\rangle^{\frac{1}{r}} \\ &\quad - \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j} \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j}} \right. \\ &\quad \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

By taking the supremum on the unit vector x in \mathcal{H} we reach the desired inequality. \square

Corollary 2.7. *Let $A_i, B_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then for $r, p \geq 1$ we have*

$$w_p^p(A_1^* B_1, \dots, A_n^* B_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n (|B_i|^{2rp} + |A_i|^{2rp}) \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x),$$

where

$$\eta(x) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle |A_i|^{2p} x, x \rangle^{2^{j-1}-k_j} \langle |B_i|^{2p} x, x \rangle^{k_j}} - \sqrt[2^j]{\langle |B_i|^{2p} x, x \rangle^{k_j+1} \langle |A_i|^{2p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.$$

Proof. Choosing $f(t) = g(t) = t^{\frac{1}{2}}$ and $T_i = I$ for $i = 1, 2, \dots, n$ in Theorem 2.6, we get the desired result. \square

Corollary 2.8. Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$), and let f and g be nonnegative continuous functions on $[0, \infty)$ such that $f(t)g(t) = t$ for all $t \in [0, \infty)$ and $r, p \geq 1$. Then

$$w_p^p(T_1, \dots, T_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n (f^{2rp}(|T_i|) + g^{2rp}(|T_i^*|)) \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x), \quad (2.3)$$

where

$$\eta(x) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle g^{2p}(|T_i^*|)x, x \rangle^{2^{j-1}-k_j} \langle f^{2p}(|T_i|)x, x \rangle^{k_j}} - \sqrt[2^j]{\langle f^{2p}(|T_i|)x, x \rangle^{k_j+1} \langle g^{2p}(|T_i^*|)x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.$$

In particular,

$$w_p^p(T_1, \dots, T_n) \leq \frac{1}{2} \left\| \sum_{i=1}^n (|T_i|^{2\alpha p} + |T_i^*|^{2(1-\alpha)p}) \right\| - \inf_{\|x\|=1} \eta(x), \quad (2.4)$$

where $0 \leq \alpha \leq 1$ and

$$\eta(x) = \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.$$

Proof. Selecting $A_i = B_i = I$ for $i = 1, 2, \dots, n$ in Theorem 2.6, we get the first result. Letting $f(t) = t^\alpha$, $g(t) = t^{1-\alpha}$, $r = 1$ and $B_i = A_i = I$ for $i = 1, 2, \dots, n$ in inequality (2.3), we reach the second inequality. \square

Remark 2.9. Note that inequality (2.4) is a refinement of inequality (1.5), since

$$\frac{1}{2} \sum_{i=1}^n (\langle |T_i|^{2\alpha p} x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{\frac{1}{2}})^2 \leq \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.$$

Now, by letting $n = 2$, $N = 1$, $T_1 = B$ and $T_2 = C$ in Theorem 2.6, we obtain the following consequence.

Corollary 2.10. Let $B, C \in \mathbb{B}(\mathcal{H})$. Then, for all $p \geq 1$ and $0 \leq \alpha \leq 1$,

$$w_p^p(B, C) \leq \frac{1}{2} \|B|^{2\alpha p} + |B^*|^{2(1-\alpha)p} + |C|^{2\alpha p} + |C^*|^{2(1-\alpha)p}\| - \inf_{\|x\|=1} \eta(x),$$

where

$$\eta(x) = \frac{1}{2} [\langle |B|^{2\alpha p} x, x \rangle^{\frac{1}{2}} - \langle |B^*|^{2(1-\alpha)p} x, x \rangle^{\frac{1}{2}}]^2 + [\langle |C|^{2\alpha p} x, x \rangle^{\frac{1}{2}} - \langle |C^*|^{2(1-\alpha)p} x, x \rangle^{\frac{1}{2}}]^2].$$

Theorem 2.11. Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then

$$w_p(T_1, \dots, T_n) \leq \frac{1}{2} \left[\sum_{i=1}^n (\| |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} \| - 2 \inf_{\|x\|=1} \eta_i(x))^p \right]^{\frac{1}{p}}, \quad (2.5)$$

where $p \geq 1$, $0 \leq \alpha \leq 1$ and

$$\eta_i(x) = \frac{1}{2} \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.$$

Proof. By using of Lemma 2.1 and inequality (1.2), for any unit vector $x \in \mathcal{H}$, we have

$$\begin{aligned} \sum_{i=1}^n |\langle T_i x, x \rangle|^p &\leq \sum_{i=1}^n (\langle |T_i|^{2\alpha} x, x \rangle^{\frac{1}{2}} \langle |T_i^*|^{2(1-\alpha)} x, x \rangle^{\frac{1}{2}})^p \\ &\leq \frac{1}{2^p} \sum_{i=1}^n \left[\langle |T_i|^{2\alpha} x, x \rangle + \langle |T_i^*|^{2(1-\alpha)} x, x \rangle - \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} \right. \right. \\ &\quad \left. \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right]^p \\ &= \frac{1}{2^p} \sum_{i=1}^n \left[\langle |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} x, x \rangle - \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} \right. \right. \\ &\quad \left. \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right]^p. \end{aligned}$$

Thus,

$$\begin{aligned} \left(\sum_{i=1}^n |\langle T_i x, x \rangle|^p \right)^{\frac{1}{p}} &\leq \frac{1}{2} \left[\sum_{i=1}^n \left(\langle |T_i|^{2\alpha} x, x \rangle + \langle |T_i^*|^{2(1-\alpha)} x, x \rangle - \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j}} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j} \right. \right. \right. \\ &\quad \left. \left. \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right)^p \right]^{\frac{1}{p}} \\ &= \frac{1}{2} \left[\sum_{i=1}^n (\langle |T_i|^{2\alpha} x, x \rangle + \langle |T_i^*|^{2(1-\alpha)} x, x \rangle - 2\eta_i(x))^p \right]^{\frac{1}{p}}. \end{aligned}$$

Now, by taking the supremum over all unit vectors $x \in \mathcal{H}$, we get the desired result. \square

Remark 2.12. If $N = 1$ in inequality (2.5), then we reach inequality (1.4). It follows from

$$\begin{aligned} &\frac{1}{2} (\langle |T_i|^{2\alpha} x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)} x, x \rangle^{\frac{1}{2}})^2 \\ &\leq \frac{1}{2} \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j}} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j} - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \end{aligned}$$

that inequality (2.5) is a refinement of inequality (1.4).

Theorem 2.13. Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then, for $0 \leq \alpha \leq 1$ and $p \geq 2$,

$$w_p^p(T_1, \dots, T_n) \leq \left\| \sum_{i=1}^n (\alpha |T_i|^p + (1-\alpha) |T_i^*|^p) \right\| - \inf_{\|x\|=1} \eta(x), \quad (2.6)$$

where

$$\begin{aligned} \eta(x) &= \sum_{i=1}^n \left(\sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \alpha + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \right. \\ &\quad \times \left. \left(\sqrt[2^j]{\langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j}} \langle |T_i|^p x, x \rangle^{k_j} - \sqrt[2^j]{\langle |T_i|^p x, x \rangle^{k_j+1} \langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right) \end{aligned}$$

Proof. For every unit vector $x \in \mathcal{H}$, by Lemma 2.1 (b), Lemma 2.2 (b) and (1.2), we have

$$\begin{aligned} \sum_{i=1}^n |\langle T_i x, x \rangle|^p &= \sum_{i=1}^n (|\langle T_i x, x \rangle|^2)^{\frac{p}{2}} \\ &\leq \sum_{i=1}^n (\langle |T_i|^{2\alpha} x, x \rangle \langle |T_i^*|^{2(1-\alpha)} x, x \rangle)^{\frac{p}{2}} \\ &\leq \sum_{i=1}^n (\langle |T_i| p x, x \rangle^\alpha \langle |T_i^*|^p x, x \rangle^{1-\alpha}) \\ &\leq \sum_{i=1}^n (\alpha \langle |T_i|^p x, x \rangle + (1-\alpha) \langle |T_i^*|^p x, x \rangle) - \sum_{i=1}^n \left(\sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \alpha + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \right. \\ &\quad \times \left. \left(\sqrt[2^j]{\langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j}} \langle |T_i|^p x, x \rangle^{k_j} - \sqrt[2^j]{\langle |T_i|^p x, x \rangle^{k_j+1} \langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right) \\ &\leq \sum_{i=1}^n \langle (\alpha |T_i|^p + (1-\alpha) |T_i^*|^p) x, x \rangle - \inf_{\|x\|=1} \eta(x) \\ &= \left\langle \sum_{i=1}^n (\alpha |T_i|^p + (1-\alpha) |T_i^*|^p) x, x \right\rangle - \inf_{\|x\|=1} \eta(x). \end{aligned}$$

Now by taking the supremum over a unit vector $x \in \mathcal{H}$ we get the desired result. \square

Remark 2.14. If we put $N = 1$ in inequality (2.6), then we get inequality (1.6). Hence, inequality (2.6) is a refinement of (1.6).

In [12, Remark 3.10], Moslehian et al. showed

$$w_p^p(B, C) \leq \frac{1}{2} \| |B|^p + |B^*|^p + |C|^p + |C^*|^p \|, \quad (2.7)$$

in which, $B, C \in \mathbb{B}(\mathcal{H})$ and $p \geq 2$. In the following result we show a refinement of (2.7).

Corollary 2.15. *Let $B, C \in \mathbb{B}(\mathcal{H})$. Then, for $p \geq 2$,*

$$w_p^p(B, C) \leq \frac{1}{2} \| |B|^p + |B^*|^p + |C|^p + |C^*|^p \| - \inf_{\|x\|=1} \eta(x), \quad (2.8)$$

where $\eta(x) = \frac{1}{2} ((\langle |B|^p x, x \rangle^{\frac{1}{2}} - \langle |B^*|^p x, x \rangle^{\frac{1}{2}})^2 - (\langle |C|^p x, x \rangle^{\frac{1}{2}} - \langle |C^*|^p x, x \rangle^{\frac{1}{2}})^2)$.

In particular, if $A \in \mathbb{B}(\mathcal{H})$, then

$$w^2(A) \leq \frac{1}{2} \| A^* A + A A^* \|.$$

Proof. If we take $N = 1$, $n = 2$, $T_1 = B$, $T_2 = C$, and $\alpha = \frac{1}{2}$ in Theorem 2.13, we get the first inequality.

In this particular case, let $A = B + iC$ be the Cartesian decomposition of A . Then $A^* A + A A^* = 2(B^2 + C^2)$ and $\inf_{\|x\|=1} \eta(x) = 0$. Thus, for $p = 2$, inequality (2.8) can be written as

$$w_2^2(B, C) \leq \| B^2 + C^2 \| = \frac{1}{2} \| A^* A + A A^* \|.$$

The desired inequality follows by noting that

$$w_2^2(B, C) = \sup_{\|x\|=1} \{ |\langle Bx, x \rangle|^2 + |\langle Cx, x \rangle|^2 \} = \sup_{\|x\|=1} |\langle Ax, x \rangle|^2 = w^2(A). \quad \square$$

Theorem 2.16. *Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$), $r \geq 1$, and $p \geq q \geq 1$, with $\frac{1}{p} + \frac{1}{q} = \frac{1}{r}$. Then*

$$w_p^r(|T_1|, \dots, |T_n|) w_q^r(|T_1^*|, \dots, |T_n^*|) \leq \frac{r}{p} \left\| \sum_{i=1}^n |T_i|^p \right\| + \frac{r}{q} \left\| \sum_{i=1}^n |T_i^*|^q \right\| - \inf_{\|x\|=\|y\|=1} \lambda(x, y), \quad (2.9)$$

where

$$\begin{aligned} \lambda(x, y) &= \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \left(\sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j} } \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j} \right. \\ &\quad \left. - \sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j-1} } \right)^2. \end{aligned}$$

Proof. Let $x, y \in \mathbb{B}(\mathcal{H})$ be unit vectors. Applying inequality (1.2) and Lemma (2.2) (a), we get

$$\begin{aligned} &\left(\left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{\frac{1}{q}} \right)^r \\ &\leq \frac{r}{p} \sum_{i=1}^n \langle |T_i| x, y \rangle^p + \frac{r}{q} \sum_{i=1}^n \langle |T_i^*| x, y \rangle^q - \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\ &\quad \times \left(\sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j} } \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j} - \sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j-1} } \right)^2 \\ &\leq \frac{r}{p} \sum_{i=1}^n \langle |T_i|^p x, y \rangle + \frac{r}{q} \sum_{i=1}^n \langle |T_i^*|^q x, y \rangle - \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\ &\quad \times \left(\sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j} } \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j} - \sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j-1} } \right)^2 \\ &= \frac{r}{p} \left\langle \left(\sum_{i=1}^n |T_i|^p \right) x, y \right\rangle + \frac{r}{q} \left\langle \left(\sum_{i=1}^n |T_i^*|^q \right) x, y \right\rangle - \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\ &\quad \times \left(\sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j} } \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j} - \sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i| x, y \rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*| x, y \rangle^q \right)^{2^{j-1}-k_j-1} } \right)^2. \end{aligned}$$

By taking the supremum on $x, y \in \mathcal{H}$ with $\|x\| = \|y\| = 1$, we get the desired inequality. \square

Remark 2.17. If we put $N = 1$ in inequality (2.9), then we come to inequality (1.7). It follows from

$$\begin{aligned} & \frac{r}{p} \left(\sqrt[2^j]{\sum_{i=1}^n \langle |T_i|x, y \rangle^p} - \sqrt[2^j]{\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q} \right)^2 \\ & \leq \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \left(\sqrt[2^j]{\left(\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q \right)^{2^{j-1}-k_j}} \left(\sum_{i=1}^n \langle |T_i|x, y \rangle^p \right)^{k_j} \right. \\ & \quad \left. - \sqrt[2^j]{\left(\sum_{i=1}^n \langle |T_i|x, y \rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q \right)^{2^{j-1}-k_j-1}} \right)^2 \end{aligned}$$

that inequality (2.9) is a refinement of (1.7).

Corollary 2.18. Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then

$$w_e(|T_1|, \dots, |T_n|) w_e(|T_1^*|, \dots, |T_n^*|) \leq \frac{1}{2} \left(\left\| \sum_{i=1}^n T_i^* T_i \right\| + \left\| \sum_{i=1}^n T_i T_i^* \right\| \right) - \inf_{\|x\|=\|y\|=1} \lambda(x, y),$$

where

$$\begin{aligned} \lambda(x, y) &= \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{1}{2} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\ &\quad \times \left(\sqrt[2^j]{\left(\sum_{i=1}^n \langle |T_i^*|x, y \rangle \right)^{2(2^{j-1}-k_j)} \left(\sum_{i=1}^n \langle |T_i|x, y \rangle \right)^{2k_j}} \right. \\ & \quad \left. - \sqrt[2^j]{\left(\sum_{i=1}^n \langle |T_i|x, y \rangle \right)^{2(k_j+1)} \left(\sum_{i=1}^n \langle |T_i^*|x, y \rangle \right)^{2(2^{j-1}-k_j-1)}} \right)^2. \end{aligned}$$

Proof. The result is obtained by letting $p = q = 2$ and $r = 1$ in inequality (2.9). \square

Corollary 2.19. Let $T_1, \dots, T_n \in \mathbb{B}(\mathcal{H})$ be positive operators. Then

$$w_e(T_1, \dots, T_n) \leq \left\| \sum_{i=1}^n T_i^2 \right\|^{\frac{1}{2}}.$$

References

- [1] O. Axelsson, H. Lu and B. Polman, On the numerical radius of matrices and its application to iterative solution methods, *Linear Multilinear Algebra* **37** (1994), no. 1–3, 225–238.
- [2] M. Bakherad, M. Krnić and M. S. Moslehian, Reverse Young-type inequalities for matrices and operators, *Rocky Mountain J. Math.* **46** (2016), no. 4, 1089–1105.
- [3] M. Boumazgour and H. A. Nabwey, A note concerning the numerical range of a basic elementary operator, *Ann. Funct. Anal.* **7** (2016), no. 3, 434–441.
- [4] K. E. Gustafson and D. K. M. Rao, *Numerical Range*, Universitext, Springer, New York, 1997.
- [5] M. Hajmohamadi, R. Lashkaripour and M. Bakherad, Some generalizations of numerical radius on off-diagonal part of 2×2 operator matrices, *J. Math. Inequal.* **12** (2018), no. 2, 447–457.
- [6] P. R. Halmos, *A Hilbert Space Problem Book*, 2nd ed., Grad. Texts in Math. 19, Springer, New York, 1982.
- [7] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Cambridge Math. Lib., Cambridge University, Cambridge, 1988.
- [8] R. Kaur, M. S. Moslehian, M. Singh and C. Conde, Further refinements of the Heinz inequality, *Linear Algebra Appl.* **447** (2014), 26–37.
- [9] F. Kittaneh, Notes on some inequalities for Hilbert space operators, *Publ. Res. Inst. Math. Sci.* **24** (1988), no. 2, 283–293.
- [10] F. Kittaneh and Y. Manasrah, Improved Young and Heinz inequalities for matrices, *J. Math. Anal. Appl.* **361** (2010), no. 1, 262–269.
- [11] M. S. Moslehian and M. Sattari, Inequalities for operator space numerical radius of 2×2 block matrices, *J. Math. Phys.* **57** (2016), no. 1, Article ID 015201.

- [12] M. S. Moslehian, M. Sattari and K. Shebrawi, Extensions of Euclidean operator radius inequalities, *Math. Scand.* **120** (2017), no. 1, 129–144.
- [13] G. Popescu, Unitary invariants in multivariable operator theory, *Mem. Amer. Math. Soc.* **200** (2009), no. 941, 1–91.
- [14] M. Sababheh and D. Choi, A complete refinement of Young's inequality, *J. Math. Anal. Appl.* **440** (2016), no. 1, 379–393.
- [15] M. Sattari, M. S. Moslehian and T. Yamazaki, Some generalized numerical radius inequalities for Hilbert space operators, *Linear Algebra Appl.* **470** (2015), 216–227.
- [16] A. Sheikholesini, M. S. Moslehian and K. Shebrawi, Inequalities for generalized Euclidean operator radius via Young's inequality, *J. Math. Anal. Appl.* **445** (2017), no. 2, 1516–1529.
- [17] T. Yamazaki, On upper and lower bounds for the numerical radius and an equality condition, *Studia Math.* **178** (2007), no. 1, 83–89.
- [18] A. Zamani, Some lower bounds for the numerical radius of Hilbert space operators, *Adv. Oper. Theory* **2** (2017), no. 2, 98–107.